arXiv:1112.0642v1 [math.CO] 3 Dec 2011

CLASSIFICATION OF INDECOMPOSABLE FLOWS OF SIGNED GRAPHS

BEIFANG CHEN AND JUE WANG

Abstract. An indecomposable flow f on a signed graph Σ is a nontrivial integral flow that cannot be decomposed into $f = f_1 + f_2$, where f_1, f_2 are nontrivial integral flows having the same sign (both ≥ 0 or both ≤ 0) at each edge of Σ. This paper is to classify indecomposable flows into characteristic vectors of circuits and Eulerian cycle-trees — a class of signed graphs having a kind of tree structure in which all cycles can be viewed as vertices of a tree. Moreover, each indecomposable flow other than circuit characteristic vectors can be further decomposed into a sum of certain half circuit characteristic vectors having the same sign at each edge. The variety of indecomposable flows of signed graphs is much richer than that of ordinary unsigned graphs.

1. Introduction

A signed graph is an ordinary graph whose each edge is endowed with either a positive sign $+1$ or a negative sign -1. The system was first introduced by Heider [9] to study social psychological relation in social balance theory, where positive and negative edges represent positive and negative relations holding between two individuals respectively. Later, Harary [8] formally introduced signed graphs and characterized balanced signed graphs up to switching — the existence of bipartition of the vertex set so that all edges inside each part are positive and all edges between the two parts are negative. After that Zaslavsky [14, 15] introduced notions of ordinary graphs to signed graphs such as circuit, bond, orientation, incidence matrix, Laplace matrix, and associated matroids, etc. With such notions, Chen and Wang [5] further introduced flow and tension lattices of signed graphs and obtained fundamental properties on flows and tensions, including a few characterizations of cuts and bonds. For the flow and tension lattices, it is natural to ask how integral flows and tensions are build up from more basic integral flows and tensions. What does an integral flow or tension look like if it cannot be further decomposed?

For ordinary graphs, it is easy to see that indecomposable flows are simply the characteristic vectors of directed circuits. For signed graphs, characteristic vectors of directed circuits are certainly indecomposable flows; however, we shall see that the indecomposable flows other than circuit characteristic vectors consist of an additional class of characteristic vectors of so-called directed Eulerian cycle-trees at integer scale. The present paper is to classify this additional class of characteristic vectors, to study the structures of indecomposable flows, and to decompose

---

2000 *Mathematics Subject Classification.* 05C20, 05C21, 05C22, 05C38, 05C45, 05C75.

Key words and phrases. Signed graph, orientation, circuit, indecomposable flow, directed Eulerian walk, minimal directed Eulerian walk.

Research is supported by RGC Competitive Earmarked Research Grants 600608 and 600409.

indecomposable flows further into characteristic vectors of directed circuits at half-integer scale. The classification is crucial to studying geometry of the flow lattices of signed graphs.

Let $\Sigma = (V, E, \sigma)$ be a signed graph throughout, where (V, E) is an ordinary finite graph with possible loops and multiple edges, V is the vertex set, E is the edge set, and $\sigma : E \to \{-1, 1\}$ is the sign function. For each edge subset $S \subseteq E$, we define the induced *signed subgraph* $\Sigma(S) := (V, S, \sigma|_S)$, where $\sigma|_S$ is the restriction of σ to S.

Recall that a *cycle* of Σ is a simple closed path. The *sign* of a cycle is the product of signs on its edges. A cycle is said to be *balanced (unbalanced)* if its sign is positive (negative). A signed graph is said to be *balanced* if all cycles are balanced, and *unbalanced* if one of its cycles is unbalanced. A connected component of Σ is called a *balanced (unbalanced) component* if it is balanced (unbalanced) as a signed subgraph.

An *orientation* of a signed graph Σ is an assignment that each edge e is assigned two arrows at its end-vertices u, v as follows: (i) if e is a positive edge, the two arrows are in the same direction; (ii) if e is a negative edge, the two arrows are in opposite directions; see the following demonstration.



We may think of an arrow on the edge e at its one end-vertex u as $+1$ if the arrow points away from u and -1 if the arrow points towards u. Then there are both $+1$ and -1 for a positive loop at its unique end-vertex, and two $+1$'s or two -1's for a negative loop at its unique end-vertex. So an orientation on Σ can be considered as a multi-valued function $\varepsilon : V \times E \to \{-1, 0, 1\}$ such that (i) $\varepsilon(u, x)$ has the double-value ± 1 or ∓ 1 if x is a positive loop at its unique end-vertex u, and is single-valued otherwise; (ii) $\varepsilon(u, x) = 0$ if u is not an end-vertex of x; and (iii)

$$\varepsilon(u, x)\varepsilon(v, x) = -\sigma(x), \quad x = uv. \tag{1.1}$$

A signed graph Σ with an orientation ε is referred to an *oriented signed graph* (Σ, ε).

Let A be an abelian group and be assumed automatically a $\mathbb{Z}$-module. Associated with an orientation ε is the *boundary operator* $\partial_\varepsilon : A^E \to A^V$ defined by

$$(\partial_\varepsilon f)(v) = \sum_{e \in E} \boldsymbol{m}_{v,e} f(e),$$

where $f \in A^E$, $v \in V$, and

$$\boldsymbol{m}_{v,e} = \begin{cases} \varepsilon(v, e) & \text{if } e \text{ is a non-loop,} \\ 2\varepsilon(v, e) & \text{if } e \text{ is a negative loop,} \\ 0 & \text{otherwise.} \end{cases} \tag{1.2}$$

A function $f : E \to A$ is said to be *conservative* at a vertex v if $(\partial_\varepsilon f)(v) = 0$, and is called an *$A$-flow* of (Σ, ε) if it is conservative at every vertex. The *support* of a flow f is the edge subset

$$\operatorname{supp} f = \{e \in E \mid f(e) \neq 0\}. \tag{1.3}$$

The set of all A-flows forms an abelian group, called the *flow group* of (Σ, ε) with values in A, denoted $F(\Sigma, \varepsilon; A)$. We refer the vector space $F(\Sigma, \varepsilon) := F(\Sigma, \varepsilon; \mathbb{R})$ to *flow space*, and the lattice $Z(\Sigma, \varepsilon) := F(\Sigma, \varepsilon; \mathbb{Z})$ to *flow lattice* of (Σ, ε). For further information about flows of signed graphs, see [1, 4, 5, 6, 10]. For notions of graphs, we refer to the books [2, 3, 7]

We call a flow to be *nontrivial* if its support is nonempty. A nontrivial integral flow f is said to be *decomposable* if f can be written as $f = f_1 + f_2$, where f_i are nontrivial integral flows having the same sign, that is, $f_1(e)f_2(e) \geq 0$ for all $e \in E$. Nontrivial integral flows that are not decomposable are referred to *indecomposable flows*. An integral flow f is said to be *elementary* if it is indecomposable and there is no nontrivial flow g such that $\operatorname{supp} g$ is properly contained in $\operatorname{supp} f$; see Tutte's definition of elementary chains [12, 13].

Let W be a walk of length n in Σ and be written as a vertex-edge sequence

$$W = u_0 x_1 u_1 x_2 \ldots u_{n-1} x_n u_n, \tag{1.4}$$

where each x_i is an edge with end-vertices u_{i-1}, u_i. The walk W is said to be *closed* if the *initial vertex* u_0 is the same as the *terminal vertex* u_n. The *support* of W is a signed subgraph $\Sigma(W)$ induced by the vertex and edge subsets

$$V(W) = \{u_0, u_1, \ldots, u_n\}, \quad E(W) = \{x_1, x_2, \ldots, x_n\}.$$

A *direction* of W is a function ε_W with values either 1 or -1, defined for all pairs (u_{i-1}, x_i) and (u_i, x_i), such that

$$\varepsilon_W(u_{i-1}, x_i)\varepsilon_W(u_i, x_i) = -\sigma(x_i), \quad \varepsilon_W(u_i, x_i) + \varepsilon_W(u_i, x_{i+1}) = 0.$$

Every walk has exactly two opposite directions. A walk W with a direction ε_W is called a *directed walk*, denoted (W, ε_W). If in addition $u_0 = u_n$ and $\varepsilon_W(u_0, x_1) + \varepsilon_W(u_n, x_n) = 0$, the directed walk (W, ε_W) is called a *closed directed walk*. A *directed Eulerian walk* is a directed closed walk whose orientations on each repeated edge are the same. We view each directed walk (W, ε_W) as a multiset

$$M(W, \varepsilon_W) = \{(x_1, \varepsilon_W(u_1, x_1)), \ldots, (x_1, \varepsilon_W(u_l, x_l))\}$$

of oriented edges. A directed Eulerian walk (W, ε_W) is said to be *minimal* if there is no directed Eulerian walk $(W', \varepsilon_{W'})$ such that $M(W', \varepsilon_{W'})$ is a proper sub-multiset of $M(W, \varepsilon_W)$, and to be *elementary* if it is minimal and there is no directed Eulerian walk $(W', \varepsilon_{W'})$ such that $E(W')$ is a proper subset of $E(W)$. A closed walk W is said to be an Eulerian walk of (Σ, ε) if (W, ε) is a directed Eulerian walk.

Let ε_i $(i = 1, 2)$ be orientations on signed subgraphs Σ_i of Σ. The *coupling* of $\varepsilon_1, \varepsilon_2$ is a function $[\varepsilon_1, \varepsilon_2] : E \to \mathbb{Z}$, defined for $x = uv$ by

$$[\varepsilon_1, \varepsilon_2](x) = \begin{cases} 1 & \text{if } x \in \Sigma_1 \cap \Sigma_2,\ \varepsilon_1(u, x) = \varepsilon_2(u, x), \\ -1 & \text{if } x \in \Sigma_1 \cap \Sigma_2,\ \varepsilon_1(u, x) \neq \varepsilon_2(u, x), \\ 0 & \text{otherwise.} \end{cases} \tag{1.5}$$

In other words, $[\varepsilon_1, \varepsilon_2](x) = \varepsilon_1(u, x)\varepsilon_2(u, x)$ if $x = uv$.

Let (W, ε_W) be a directed closed walk of length n with the vertex-edge sequence (1.4), where $x_i = u_{i-1}u_i$. We may think of the walk W as an edge multiset $\{x_1, x_2, \ldots, x_n\}$ of n elements. The *characteristic vector* of (W, ε_W) on (Σ, ε) is a function $f_W : E \to \mathbb{Z}$ defined by

$$f_W(x) = \sum_{x_i \in W,\, x_i = x} [\varepsilon, \varepsilon_W](x_i). \tag{1.6}$$

It can be shown that f_W is an integral flow of (Σ, ε); see Lemma 2.1.

Theorem 1.1 (Flow Reduction)**.** *Let f be a nontrivial integral flow of an oriented signed graph (Σ, ε). Then there exist minimal directed Eulerian walks (W_i, ε_{W_i}) such that*

$$f = \sum f_{W_i},$$

where f_{W_i} are defined by (1.6)*. Furthermore, if f is indecomposable, then $f = f_W$ for a minimal directed Eulerian walk (W, ε_W).*

A *cycle-tree* of Σ is a connected signed subgraph T which can be decomposed into edge-disjoint cycles C_i and internally vertex-disjoint simple paths P_j (called *tree-paths*), denoted $T = \{C_i, P_j\}$, satisfying the four conditions: (i) there is no cycle other than C_i in T; (ii) the intersection of two cycles is either empty or a single vertex (called an *intersection vertex*); (iii) each P_j intersects exactly two cycles and the intersections are exactly the initial and terminal vertices (also called *intersection vertices*); and (iv) each intersection vertex is a *cut-point* (a vertex whose removal increases the number of connected components of the underline graph (as a topological space of 1-dimensional CW complex), also called a *separating vertex* (see [3], p.119). A cycle-tree is said to be *Eulerian* if it further satisfies (v) *Parity Condition*: each balanced cycle has even number of intersection vertices and each unbalanced cycle has odd number of intersection vertices.

We call a cycle in a cycle-tree to be an *end-cycle* if it has exactly one intersection vertex. The name cycle-tree is justified as follows: if one converts each cycle C_i into a vertex, each tree-path P_j into an edge adjacent with the two vertices converted from the two cycles connected by P_j, and each (possible) nonempty intersection vertex of two cycles into an edge adjacent with the two vertices converted from the two cycles, then the graph obtained by the conversion is indeed a tree.

We shall see that each Eulerian cycle-tree can be endowed with an orientation such that every vertex is neither a sink nor a source; such an orientation is called a *direction* of the Eulerian cycle-tree. We shall further see that there are exactly two directions on each Eulerian cycle-tree, and these two directions are opposite each other. An Eulerian cycle-tree T with a direction ε_T is referred to a *directed Eulerian cycle-tree* (T, ε_T).

Let $T = \{C_i, P_j\}$ be an Eulerian cycle-tree of Σ. The *indicator* of T is a function $I_T : E \to \{0, 1, 2\}$ defined by

$$I_T(x) = \begin{cases} 1 & \text{if } x \text{ is on a cycle of } T, \\ 2 & \text{if } x \text{ is on a tree-path of } T, \\ 0 & \text{otherwise.} \end{cases} \tag{1.7}$$

Given a direction ε_T of T. Viewing both (Σ, ε) and (T, ε_T) as oriented signed subgraphs of Σ, we have the coupling $[\varepsilon, \varepsilon_T]$. The product function $[\varepsilon, \varepsilon_T]I_T$ determines a vector in $\mathbb{Z}^E$ and is an integral flow of (Σ, ε), called the *characteristic vector* of the directed Eulerian cycle-tree (T, ε_T) for (Σ, ε).

Theorem 1.2 (Realization of Minimal Directed Eulerian Walk)**.** *Let $T = \{C_i, P_j\}$ be a cycle-tree. Then there exists a closed walk W so that the edges on the cycles C_i appear exactly once and the edges on the tree-paths P_j appear exactly twice.*

Moreover, if T is further Eulerian, then each direction ε_T of T gives automatically a direction of W so that (W, ε_T) is a minimal directed Eulerian walk.

An Eulerian cycle-tree T is called a *circuit* if T is minimal in the sense that T does not contain properly any Eulerian cycle-tree. It can be easily seen that each circuit C must be one of the following three types: (i) consisting of a single balanced cycle, said to be of *Type I*; (ii) consisting of two unbalanced cycles C_1, C_2 whose intersection is a single vertex, written $C = C_1C_2$ and said to be of *Type II*; (iii) consisting of two vertex-disjoint unbalanced cycles C_1, C_2, and a simple path P having at least one edge, written $C = C_1PC_2$ and said to be of *Type III*, such that $C_1 \cap P$ is the initial vertex and $C_2 \cap P$ the terminal vertex of P. The present definition of circuit is equivalent to that defined in [14] and that adopted in [5, 6].

Theorem 1.3 (Characterization of Minimal Directed Eulerian Walk). *Let* (W, ε_W) *be a minimal directed Eulerian walk. Then*

(a) The support of W *is an Eulerian cycle-tree* T_W*, called the* supporting Eulerian cycle-tree *of* W*; edges on the cycles of* T_W *appear exactly once in* W*, while edges on the tree-paths of* T_W *appear exactly twice in* W*. Moreover, the direction* ε_W *induces a direction on* T_W*.*

(b) If T_W *is a circuit, then (i)* $W = C$*, a balanced cycle; (ii)* $W = C_1C_2$*, where* C_1, C_2 *are unbalanced cycles whose intersection is a single vertex; (iii)* $W = C_1PC_2P^{-1}$*, where* C_1, C_2 *are vertex-disjoint unbalanced cycles, and* P *is a simple path intersecting* C_1 *exactly at its initial vertex and* C_2 *exactly at its terminal vertex.*

(c) If T_W *is not a circuit, then the vertex-edge sequence of* W *can be divided into the form*

$$W = C_1P_1C_2P_2 \cdots C_kP_kC_{k+1}, \quad k \geq 2,$$

where C_i *are vertex-disjoint unbalanced cycles with* $C_{k+1} = C_1$*,* P_i *are simple open paths having possible double vertices and double edges, and each* $C_iP_iC_{i+1}$ *is a circuit of Type III. Furthermore, the function* f_W *(defined by (1.6)) is an indecomposable flow of* (Σ, ε) *and*

$$f_W = \frac{1}{2}\sum_{i=1}^{k}[\varepsilon, \varepsilon_W]I_{C_iP_iC_{i+1}}. \tag{1.8}$$

(d) The orientations of edges at a double vertex of W *have the patterns demonstrated in Figure 1. Moreover, if a path* P_s *intersects the paths* P_r *with* $r < s$ *at the vertices* $u_{\gamma_1}, u_{\gamma_2}, \ldots, u_{\gamma_m}$ *with index order* $\gamma_1 < \gamma_2 < \cdots < \gamma_m$ *in* P_s*, then the same vertices* $u_{\beta_1}, u_{\beta_2}, \ldots, u_{\beta_m}$ *have the index order* $\beta_1 > \beta_2 > \cdots > \beta_m$ *in the paths* P_r*.*



FIGURE 1. Orientation patterns at a double vertex.

Corollary 1.4. *Let* f *be a nontrivial integral flow of an oriented signed graph* (Σ, ε)*. Then* f *is indecomposable if and only if* f *is the characteristic vector of a directed Eulerian cycle-tree* (T, ε_T)*, that is,*

$$f = [\varepsilon, \varepsilon_T]I_T.$$

Corollary 1.5. *Let f be a nontrivial integral flow f of an oriented signed graph (Σ, ε). Then the following three statements are equivalent.*

- (a) *The flow f is elementary.*
- (b) *The flow f is the characteristic vector of a directed circuit.*
- (c) *There exists an elementary minimal directed Eulerian walk (W, ε_W) such that $f = f_W$.*

Remark. Corollary 1.5 was obtained by Bouchet [4] (see Corollary 2.3, p.283), using Zaslavsky's characterization of circuits [14]. As Zaslavsky pointed out himself, the central observation of [14] (p.53) is the existence of a matroid over the edge set of a signed graph whose circuits are exactly those of Types I, II, III. Bouchet [4] assumed (without argument) that this matroid is the same as the matroid whose circuits are the supports of elementary flows. Indeed, it is trivial to see that the circuits of the formal are the circuits of the latter. However, the converse seems not straightforward without argument, though it is anticipated. Theorem 1.3 implies immediately that the converse is indeed also true. Now it is logically clear and complete that the matroid constructed by Zaslavsky [14] for a signed graph is the same as the matroid whose circuits are the supports of elementary chains (=elementary flows) of the signed graph in the sense of Tutte [13]; so are their dual matroids.

2. Flow Reduction Algorithm

Recall that a nontrivial integral flow f of a signed digraph (Σ, ε) is *decomposable* if there exist nontrivial integral flows f_1, f_2 such that $f = f_1 + f_2$ and $f_1(e)f_2(e) \geq 0$ for all $e \in E$, that is, f_1, f_2 have the same sign (both nonnegative or both nonpositive) at every edge. If follows that if $f(e) = 0$ then $f_1(e) = f_2(e) = 0$, and that f, f_i have the same sign at every edge.

Given a nonzero integer-valued function $f \in \mathbb{Z}^E$. Let ε_f be an orientation on Σ defined by

$$\varepsilon_f(u, x) = \begin{cases} -\varepsilon(u, x) & \text{if } f(e) < 0,\ x = uv, \\ \varepsilon(u, x) & \text{otherwise.} \end{cases} \tag{2.1}$$

It is trivial to see that f is a flow of (Σ, ε) if and only if the absolute value function $|f|$ is a flow of (Σ, ε_f). Moreover, if f is decomposable into $f = f_1 + f_2$ with $f_1(e)f_2(e) \geq 0$, then $|f|$ is decomposable into $|f| = |f_1| + |f_2|$, and vice versa. So, to decompose a flow f of (Σ, ε), it is equivalent to decompose the nonnegative flow $|f|$ of (Σ, ε_f). Thus without loss of generality, we only consider nontrivial nonnegative integral flows of (Σ, ε).

Now a nonnegative, nontrivial, integral flow f on (Σ, ε) is *indecomposable* provided that, if f_1 is a nonnegative, nontrivial, integral flow of (Σ, ε) and $f_1 \leq f$, then $f_1 = f$. The following *Flow Reduction Algorithm* (FRA) finds a minimal directed Eulerian walk in explicit pattern from a given nonnegative nontrivial integral flow. Let us first show that the characteristic vector of a directed closed walk is a flow.

Lemma 2.1. *Let (W, ε_W) be a directed closed walk. Then the function f_W defined by* (1.6) *is an integral flow of* (Σ, ε).

Proof. Let $W = u_0 x_1 u_1 x_2 \ldots x_n u_n$ be the vertex-edge sequence, where each edge x_i has end-vertices u_{i-1}, u_i, and $u_n = u_0$. Fix a vertex $v \in V$. Let E_v be the set of edges incident with v, and let $u_{a_1}, u_{a_2}, \ldots, u_{a_k}$ be the sequence that v appears in

W. Since $\varepsilon_W(u_{a_i},x_{a_i})+\varepsilon_W(u_{a_i},x_{a_i+1})=0$, we have

$$\begin{aligned}\sum_{x\in E_v}\varepsilon(v,x)f_W(x) &= \sum_{x\in E_v}\sum_{x_i\in W,\, x_i=x}\varepsilon(v,x_i)[\varepsilon,\varepsilon_W](x_i)\\ &= \sum_{x_i\in W,\, x_i\in E_v}\varepsilon_W(v,x_i)\\ &= \sum_{i=1}^{k}\big[\varepsilon_W(u_{a_i},x_{a_i})+\varepsilon_W(u_{a_i},x_{a_i+1})\big]=0.\end{aligned}$$

Hence the function f_W is an integral flow of (Σ,ε). □

Flow Reduction Algorithm (FRA)

STEP 0. Choose an edge $x_1=u_0u_1\in\operatorname{supp} f$ with end-vertices u_0,u_1. Initiate a walk $W=u_0x_1$, an *edge variable* $\boldsymbol{x}$, a *vertex variable* $\boldsymbol{u}$, and an *index function* $\ell(\boldsymbol{x})$. Initially, assign x_1 to $\boldsymbol{x}$ and set $\ell(\boldsymbol{x}):=1$; assign an end-vertex of x_1 (other than u_0 if x_1 is not a loop) to $\boldsymbol{u}$. Assume that we have obtained a directed walk in the form

$$W=u_0x_1u_1x_2\ldots x_{l-1}u_{l-1}x_l,\quad \boldsymbol{x}=x_l,\quad \boldsymbol{u}=u_l,\quad \ell(\boldsymbol{x})=l\geq 1,$$

where the vertices $u_0,u_1,\ldots,u_{l-1}$ are distinct, and assume that the function f_W is given by (1.6). Note that (W,ε) is a directed closed walk if and only if f_W is a flow.

If f_W is not a flow, so is $f'_W:=f-f_W$. The idea is to let W grow from f'_W by adding a vertex-edge segment u_lx' to the end of W, where x' is an edge in $\operatorname{supp} f'_W$.

STEP 1. The walk W is a directed path without the terminal vertex u_l, having all vertices and edges distinct; $(\boldsymbol{x},\boldsymbol{u})=(x_l,u_l)$, $l=\ell(\boldsymbol{x})\geq 1$. The vertex assigned to the variable $\boldsymbol{u}$ may or may not be a vertex in $V(W)$.

CASE 1: $\boldsymbol{u}\notin V(W)$.

Then f_W is clearly not conservative at $\boldsymbol{u}$. There exists an edge $x'\in\operatorname{supp} f'_W$ with end-vertices $\boldsymbol{u},u'$ such that $\varepsilon(\boldsymbol{u},\boldsymbol{x})\neq\varepsilon(\boldsymbol{u},x')$. Define $\ell(x')=l+1$, write $\boldsymbol{u}x'$ as u_lx_{l+1}, and set $W:=Wu_lx'$. Assign (x',u') to $(\boldsymbol{x},\boldsymbol{u})$ and return to STEP 1.

CASE 2: $\boldsymbol{u}\in V(W)$.

Let i_1 be the index of the vertex $\boldsymbol{u}$ in W and set $j_1:=l$. Note that u_l is not yet appeared in W. Then $0\leq i_1<l$ and $\boldsymbol{u}=u_{j_1}=u_{i_1}$. Define

$$\begin{aligned}C_1&=u_{i_1}x_{i_1+1}u_{i_1+1}x_{i_1+2}\ldots x_{j-1}u_{j_1-1}x_{j_1}u_{j_1},\\ Q_1&=x_{j_1+1}u_{j_1+1}x_{j_1+2}u_{j_1+2}\ldots u_{l-2}x_{l-1}u_{l-1}x_l.\end{aligned}$$

Without loss of generality, we may assume $i_1=0$, for we may select u_{i_1} as the initial vertex u_0. We thus have $\boldsymbol{u}=u_l=u_{j_1}=u_{i_1}=u_0$.

If $\varepsilon(\boldsymbol{u},\boldsymbol{x})\neq\varepsilon(u_{i_1},x_{i_1+1})$, then (C_1,ε) is a directed circuit of Type I. STOP; a directed Eulerian walk $W_0:=C_1$ is found. See the left of Figure 2.

If $\varepsilon(\boldsymbol{u},\boldsymbol{x})=\varepsilon(u_{i_1},x_{i_1+1})$, then C_1 is an unbalanced cycle, and f_W is not conservative at $\boldsymbol{u}$. Thus there exists an edge $x'=\boldsymbol{u}u'\in\operatorname{supp} f'_W$ such that $\varepsilon(\boldsymbol{u},\boldsymbol{x})\neq\varepsilon(\boldsymbol{u},x')$; see the right of Figure 2. Define $\ell(x')=l+1$, write $\boldsymbol{u}x'$ as u_lx_{l+1}, and set $W:=Wu_lx'$. Assign (x',u') to $(\boldsymbol{x},\boldsymbol{u})$; go to STEP 2.



FIGURE 2. Left: $\varepsilon(\boldsymbol{u},\boldsymbol{x}) \neq \varepsilon(u_{i_1}, x_{i_1+1})$. Right: $\varepsilon(\boldsymbol{u},\boldsymbol{x}) = \varepsilon(u_{i_1}, x_{i_1+1})$.

STEP 2. Let us write $W = C_1Q_1$, where the cycle C_1 is vertex-disjoint from the open path Q_1, $(\boldsymbol{x},\boldsymbol{u}) = (x_l, u_l)$, $l = \ell(\boldsymbol{x})$, $0 = i_1 < j_1 < l$, and $u_0 = u_{i_1} = u_{j_1}$. Note that $Q_1 = \boldsymbol{x} = x_l$ when $l = j_1 + 1$.

If $\boldsymbol{u} \notin V(W)$, it is clear that f_W is not conservative at $\boldsymbol{u}$. There exists an edge $x' = \boldsymbol{u}u' \in \operatorname{supp} f'_W$ such that $\varepsilon(\boldsymbol{u},\boldsymbol{x}) \neq \varepsilon(\boldsymbol{u}, x')$. Define $\ell(x') = l + 1$, write $\boldsymbol{u}x'$ as $u_l x_{l+1}$, and set $W := W u_l x'$. Assign (x', u') to $(\boldsymbol{x},\boldsymbol{u})$; return to STEP 2.

If $\boldsymbol{u} \in V(W)$, we have three cases.

CASE 1: $\boldsymbol{u} \in V(C_1)$ and $\boldsymbol{u} \neq u_{i_1}$.

Let α be the index of the vertex $\boldsymbol{u}$ in C_1. Then $i_1 < \alpha < j_1$. Let

$$Q'_1 = u_\alpha x_{\alpha+1} u_{\alpha+1} x_{\alpha+2} \dots u_{j_1-1} x_{j_1} u_{j_1},$$

$$Q''_1 = u_\alpha x_\alpha u_{\alpha-1} x_{\alpha-1} \dots u_{i_1+1} x_{i_1+1} u_{i_1}.$$

Note that $\varepsilon(u_{i_1}, x_{i_1+1}) = \varepsilon(u_{j_1}, x_{j_1}) \neq \varepsilon(u_{j_1}, x_{j_1+1})$, for C_1 is an unbalanced cycle.

If $\varepsilon(\boldsymbol{u},\boldsymbol{x}) \neq \varepsilon(u_\alpha, x_{\alpha+1})$, then $W_0 := Q_1Q'_1$ is a directed circuit of Type I, which is a directed Eulerian sub-walk of W. STOP; see the left of Figure 3.

If $\varepsilon(\boldsymbol{u},\boldsymbol{x}) = \varepsilon(u_\alpha, x_{\alpha+1})$, then $W_0 := Q_1Q''_1$ is a directed circuit of Type I, which is a directed Eulerian sub-walk of W. STOP; see the left of Figure 3 with the opposite orientation on $\boldsymbol{x}$.



FIGURE 3. Left: $\boldsymbol{u} = u_\alpha$. Right: $\boldsymbol{u} = u_{i_1} = u_{j_1}$.

CASE 2: $\boldsymbol{u} \in V(C_1)$ and $\boldsymbol{u} = u_{i_1}$.

Then $\boldsymbol{u} = u_{i_1} = u_{j_1} = u_l$. We have a cycle

$$C_2 = u_{j_1} x_{j_1+1} u_{j_1+1} x_{j_1+2} \dots u_{l-1} x_l u_l.$$

If $\varepsilon(\boldsymbol{u},\boldsymbol{x}) \neq \varepsilon(u_{j_1}, x_{j_1+1})$, then $W_0 := C_2$ is a directed circuit of Type I and is a directed Eulerian sub-walk of W. STOP; see the right of Figure 3.

If $\varepsilon(\boldsymbol{u},\boldsymbol{x})=\varepsilon(u_{j_1},x_{j_1+1})$, then C_2 is an unbalanced cycle. Thus $W=C_1C_2$ is a directed circuit of Type II and is a directed Eulerian walk. STOP; see the right of Figure 3 with the opposite orientation on $\boldsymbol{x}$.

CASE 3: $\boldsymbol{u}\in V(Q_1)$ and $\boldsymbol{u}\notin V(C_1)$.

Let i_2 be the index of the vertex $\boldsymbol{u}$ in Q_1 and set $j_2:=l$. Then $j_1<i_2<l$. Define

$$\begin{aligned}P_1&=x_{j_1+1}u_{j_1+1}x_{j_1+2}u_{j_1+2}\dots x_{i_2-1}u_{i_2-1}x_{i_2},\\ C_2&=u_{i_2}x_{i_2+1}u_{i_2+1}x_{i_2+2}\dots u_{j_2-1}x_{j_2}u_{j_2}.\end{aligned}$$

If $\varepsilon(\boldsymbol{u},\boldsymbol{x})\neq\varepsilon(u_{i_2},x_{i_2+1})$, then (C_2,ε) is a directed circuit of Type I. Thus $W_0:=C_2$ is a directed Eulerian sub-walk of W. STOP; see the left of Figure 4.

If $\varepsilon(\boldsymbol{u},\boldsymbol{x})=\varepsilon(u_{i_2},x_{i_2+1})$, then C_2 is an unbalanced cycle, and f_W is unbalanced at $\boldsymbol{u}$. Thus there exists an edge $x'=\boldsymbol{u}u'\in\operatorname{supp}f'_W$ such that $\varepsilon(\boldsymbol{u},\boldsymbol{x})\neq\varepsilon(\boldsymbol{u},x')$. Define $\ell(x')=l+1$, write $\boldsymbol{u}x'$ as u_lx_{l+1}, and set $W:=W\boldsymbol{u}x'$. Assign (x',u') to $(\boldsymbol{x},\boldsymbol{u})$; go to STEP 3. See the middle of Figure 4 when $x'=x_{i_2}$ and the right when $x'\neq x_{i_2}$.



FIGURE 4. Left: $\varepsilon(u_l,x_l)\neq\varepsilon(u_{i_2},x_{i_2+1})$. Middle: $\varepsilon(u_l,x_l)=\varepsilon(u_{i_2},x_{i_2+1})$, $x'=x_{i_2}$. Right: $\varepsilon(u_l,x_l)=\varepsilon(u_{i_2},x_{i_2+1})$, $x'\neq x_{i_2}$.

STEP 3. We obtain a directed walk of the form

$$W=C_1P_1C_2P_2\cdots C_{k-1}P_{k-1}C_k\boldsymbol{x},\quad k\geq 2,$$

where C_r are vertex-disjoint unbalanced cycles, P_s are simple open paths and disjoint from all C_r, $C_sP_sC_{s+1}$ are circuits of Type III, the open paths P_s may be overlapped at some vertices and edges, but no three or more are overlapped at any vertex (and subsequently at any edge). So there are only possible double vertices and possible double edges on the open paths P_s. There is no directed Eulerian sub-walk *properly* contained in W. Let us write

$$\begin{aligned}C_r&=u_{i_r}x_{i_r+1}u_{i_r+1}x_{i_r+2}\dots u_{j_r-1}x_{j_r}u_{j_r}, && 1\leq r\leq k,\\ P_s&=x_{j_s+1}u_{j_s+1}x_{j_s+2}u_{j_s+2}\dots x_{i_{s+1}-1}u_{i_{s+1}-1}x_{i_{s+1}}, && 1\leq s<k.\end{aligned}$$

If W is conservative at $\boldsymbol{u}$, then W is a directed Eulerian walk. If W is not conservative at $\boldsymbol{u}$, there exists an edge x' with end-vertices $\boldsymbol{u},u'$ such that $\varepsilon(\boldsymbol{x},\boldsymbol{u})\neq\varepsilon(\boldsymbol{u},x')$.

Adding $\boldsymbol{u}x'$ to the end of W and assign (x', u') to $(\boldsymbol{x}, \boldsymbol{u})$, we arrive at the following situation of STEP 4.

STEP 4. $W = C_1P_1C_2P_2 \cdots C_{k-1}P_{k-1}C_kQ_k$ contains no directed Eulerian sub-walks properly, where

$$Q_k = x_{j_k+1}u_{j_k+1}x_{j_k+2}u_{j_k+2} \ldots x_{l-1}u_{l-1}x_l \tag{2.2}$$

is an open path of length at least one, $l = \ell(\boldsymbol{x})$, $\boldsymbol{x} = x_l$, $\boldsymbol{u} = u_l$, $l > j_k$. In STEP 3, $l = j_k + 1$. Moreover, Q_k is vertex-disjoint from all C_r $(1 \le r \le k)$, and may intersect some P_s $(1 \le s < k)$. However, all overlapped vertices and edges are double vertices and double edges. There is no directed Eulerian sub-walk properly contained in W.

If $\boldsymbol{u} \notin V(W)$, then f_W is not conservative at $\boldsymbol{u}$. There exists an edge $x' = \boldsymbol{u}u' \in \operatorname{supp} f'_W$ such that $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) \ne \varepsilon(\boldsymbol{u}, x')$. Define $\ell(x') = l + 1$, write $\boldsymbol{u}x'$ as u_lx_{l+1}, and set $W := Wu_lx'$. Assign (x', u') to $(\boldsymbol{x}, \boldsymbol{u})$; return to STEP 4.

If $\boldsymbol{u} \in V(W)$, we divide the situation into three cases.

CASE 1: $\boldsymbol{u} \notin V(Q_k)$, $\boldsymbol{u} \in V(P_1 \cup \cdots \cup P_{k-1})$.

There are two subcases.

SUBCASE 1.1: $\boldsymbol{u} \notin V(Q_k)$, $\boldsymbol{u} \in V(P_r)$ for a unique r with $1 \le r < k$.

Let α be the index of $\boldsymbol{u}$ in P_r. Then $\boldsymbol{u} = u_\alpha$ and $j_r < \alpha < i_{r+1}$. There are two situations.

Situation 1.1.1: $V(Q_k) \cap V(P_r \cup \cdots \cup P_{k-1}) = \emptyset$.

If $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) \ne \varepsilon(u_\alpha, x_{\alpha+1})$, we define the path

$$P'_r = u_\alpha x_{\alpha+1}u_{\alpha+1}x_{\alpha+2}u_{\alpha+2} \ldots x_{i_{r+1}-1}u_{i_{r+1}-1}x_{i_{r+1}}u_{i_{r+1}}. \tag{2.3}$$

Then $W_0 := C_{r+1}P_{r+1} \cdots C_{k-1}P_{k-1}C_kQ_kP'_r$ is a directed Eulerian sub-walk of W.



FIGURE 5. $C_{r+1}P_{r+1} \cdots C_kQ_kP'_r$ is a directed Eulerian walk.

STOP; see Figure 5.

If $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) = \varepsilon(u_\alpha, x_{\alpha+1})$, then f_W is not conservative at $\boldsymbol{u}$ because $\boldsymbol{u} \ne u_{i_1} (= u_0)$. There exists an edge $x' = \boldsymbol{u}u'$ such that $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) \ne \varepsilon(\boldsymbol{u}, x')$. Define $\ell(x') = l + 1$, write $\boldsymbol{u}x'$ as u_lx_{l+1}, and set $W := Wu_lx'$. Assign (x', u') to $(\boldsymbol{x}, \boldsymbol{u})$ and return to STEP 4. See the upper one of Figure 6 when $x' = x_\alpha$ and the lower one when $x' \ne x_\alpha$.

Situation 1.1.2: $V(Q_k) \cap V(P_1 \cup \cdots \cup P_{k-1}) \ne \emptyset$.

Let γ be the greatest index for the vertices of $V(Q_k) \cap V(P_1 \cup \cdots \cup P_{k-1})$ in Q_k. Note that $\gamma < l$ because u_l is not inside Q_k. Since there is no triple vertex in W,



FIGURE 6. Upper: $x' = x_\alpha$. Lower: $x' \neq x_\alpha$.

there exists a unique s with $1 \leq s < k$ such that $u_\gamma \in V(Q_k) \cap V(P_s)$. Let β be the index of u_γ in P_s. Clearly, $u_\beta = u_\gamma$ and $\boldsymbol{u} = u_\alpha = u_l$. Since $\boldsymbol{u} \notin V(Q_k)$, it follows that $u_\alpha \neq u_\beta$, and subsequently, $\alpha \neq \beta$. (It is possible that $r = s$ and $P_r = P_s$.) We must have passed the latter step of **Situation 1.1.1** at the step $\ell(\boldsymbol{x}) = \gamma$.

If $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) \neq \varepsilon(u_\alpha, x_{\alpha+1})$, then $W_0 := C_{r+1}P_{r+1} \cdots C_k Q_k P'_r$ is a directed Eulerian sub-walk of W, where P'_r is a path defined by (2.3); STOP.

If $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) = \varepsilon(u_\alpha, x_{\alpha+1})$, there are two sub-situations.

Sub-situation 1: $\alpha < \beta$.

Then there exits an edge $x' = \boldsymbol{u}u' \in \operatorname{supp} f'_W$ such that $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) \neq \varepsilon(\boldsymbol{u}, x')$. Define $\ell(x') = l + 1$, write $\boldsymbol{u}x'$ as $u_l x_{l+1}$, and set $W := W u_l x'$. Assign (x', u') to $(\boldsymbol{x}, \boldsymbol{u})$ and return to STEP 4. See Figure 7.



FIGURE 7. $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) = \varepsilon(u_\alpha, x_{\alpha+1})$.

Sub-situation 2: $\alpha > \beta$.

Then $W_0 := C_r P'_r Q''_k P'_{r-1}$ is a directed Eulerian sub-walk of W, where

$$P'_r = (u_{j_r})x_{j_r+1}u_{j_r+1}x_{j_r+2}u_{j_r+2} \ldots x_{\alpha-1}u_{\alpha-1}x_\alpha u_\alpha,$$

$$Q''_k = (u_l)x_l u_{l-1}x_{l-1}u_{l-2} \ldots x_{\gamma+1}u_{\gamma+1}x_{\gamma+1}u_\gamma,$$

$$P'_{r-1} = (u_\beta)x_{\beta+1}u_{\beta+1}x_{\beta+2}u_{\beta+2} \ldots x_{i_r-1}u_{i_r-1}x_{i_r}u_{i_r},$$

and $u_\alpha = u_l$, $u_\gamma = u_\beta$. STOP; see Figure 8.

In this case we must have $\varepsilon(u_\gamma, x_\gamma) = \varepsilon(u_\beta, x_{\beta+1})$, for we have passed the step where $\ell(\boldsymbol{x}) = \gamma$. Moreover, if the vertices of $V(Q_k) \cap V(P_1 \cup \cdots \cup P_{k-1})$ are listed as



FIGURE 8. $C_rP_r'Q_k''P_{r-1}'$ is a directed Eulerian walk.

$u_{\gamma_1}, u_{\gamma_2}, \ldots, u_{\gamma_m}$ in Q_k with the index order $\gamma_1 < \gamma_2 < \cdots < \gamma_m$, then the same list $u_{\beta_1}, u_{\beta_2}, \ldots, u_{\beta_m}$ of the vertices with indices in $P_1, \ldots, P_{k-1}$ has the index order $\beta_1 > \beta_2 > \cdots > \beta_m$, and $\varepsilon(u_{\gamma_i}, x_{\gamma_i}) = \varepsilon(u_{\beta_i}, x_{\beta_i+1})$ for all $1 \le i \le m$.

SUBCASE 1.2: $\boldsymbol{u} \notin V(Q_k)$, $\boldsymbol{u} \in V(P_r) \cap V(P_s)$ for two indices r, s with $1 \le r < s < k$.

Let α be the index of $\boldsymbol{u}$ in P_r and β the index in P_s. Then $\boldsymbol{u} = u_\alpha = u_\beta = u_l$. Note that we must have $\varepsilon(u_\beta, x_\beta) = \varepsilon(u_\alpha, x_{\alpha+1})$.

If $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) \neq \varepsilon(u_\alpha, x_{\alpha+1})$, then $W_0 := C_{r+1}P_{r+1} \cdots C_{k-1}P_{k-1}C_kQ_kP_r'$ is a directed Eulerian sub-walk of W, where P_r' is a path defined by (2.3). STOP; see Figure 9.

If $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) = \varepsilon(u_\alpha, x_{\alpha+1})$, then $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) = \varepsilon(u_\beta, x_\beta)$. Since $\varepsilon(u_\beta, x_\beta) \neq \varepsilon(u_\beta, x_{\beta+1})$, we see that $\varepsilon(\boldsymbol{u}, \boldsymbol{x}) \neq \varepsilon(u_\beta, x_{\beta+1})$. Thus $W_0 := C_{s+1}P_{s+1} \cdots C_{k-1}P_{k-1}C_kQ_kP_s'$ is a directed Eulerian sub-walk of W, where

$$P_s' = u_\beta x_{\beta+1} u_{\beta+1} x_{\beta+2} \ldots x_{i_{s+1}-1} u_{i_{s+1}-1} x_{i_{s+1}} (u_{i_{s+1}}).$$

STOP; see Figure 9 with the opposite orientation on $\boldsymbol{x}$.



FIGURE 9. $C_{r+1}P_{r+1} \cdots C_{k-1}P_{k-1}C_kQ_kP_r'$ is a directed Eulerian walk.

CASE 2: $\boldsymbol{u} \notin V(Q_k)$, $\boldsymbol{u} \in V(C_1 \cup \cdots \cup C_k)$.

Let $\boldsymbol{u} \in V(C_r)$. Recall $C_r = u_{i_r} x_{i_r+1} u_{i_r+1} x_{i_r+2} \ldots u_{j_r-1} x_{j_r} u_{j_r}$ and $u_{i_r} = u_{j_r}$. There are two subcases.

SUBCASE 2.1: $\boldsymbol{u} \neq u_{i_r} = u_{j_r}$.

Let α be the index of $\boldsymbol{u}$ in C_r. Then $\boldsymbol{u} = u_\alpha$ and $i_r < \alpha < j_r$. Define

$$Q_r' = u_\alpha x_{\alpha+1} u_{\alpha+1} x_{\alpha+2} \ldots u_{j_r-1} x_{j_r} u_{j_r},$$

$$Q_r'' = u_\alpha x_\alpha u_{\alpha-1} x_{\alpha-1} \ldots u_{i_r+1} x_{i_r+1} u_{i_r}.$$

If $\varepsilon(\boldsymbol{u},\boldsymbol{x}) \neq \varepsilon(u_\alpha, x_{\alpha+1})$, then $W_0 := C_{r+1}P_{r+1}\cdots C_{k-1}P_{k-1}C_kQ_kQ'_rP_r$ is a directed Eulerian sub-walk of W. STOP; see Figure 10.

If $\varepsilon(\boldsymbol{u},\boldsymbol{x}) = \varepsilon(u_\alpha, x_{\alpha+1})$, then $W_0 := C_{r+1}P_{r+1}\cdots C_{k-1}P_{k-1}C_kQ_kQ''_rP_r$ is a directed Eulerian sub-walk of W. STOP; see Figure 10 with opposite orientation on $\boldsymbol{x}$.



FIGURE 10. $C_{r+1}P_{r+1}\cdots C_kQ_kQ'_rP_r$ is a directed Eulerian walk.

SUBCASE 2.2: $\boldsymbol{u} = u_{i_r} = u_{j_r}$.

If $\varepsilon(\boldsymbol{u},\boldsymbol{x}) \neq \varepsilon(u_{i_r}, x_{i_r+1})$, then $W_0 := C_rP_r\cdots C_{k-1}P_{k-1}C_kQ_k$ is a directed Eulerian sub-walk of W. STOP; see Figure 11.

If $\varepsilon(\boldsymbol{u},\boldsymbol{x}) = \varepsilon(u_{i_r}, x_{i_r+1})$, then $\varepsilon(\boldsymbol{u},\boldsymbol{x}) \neq \varepsilon(u_{j_r}, x_{j_r+1})$ because $\varepsilon(u_{i_r}, x_{i_r+1}) = \varepsilon(u_{i_r}, x_{i_r}) \neq \varepsilon(u_{j_r}, x_{j_r+1})$. Hence $W_0 := C_{r+1}P_{r+1}\cdots C_{k-1}P_{k-1}C_kQ_ku_lP_r$ is a directed Eulerian sub-walk of W. STOP; see Figure 11 with opposite orientation on $\boldsymbol{x}$.



FIGURE 11. $C_rP_r\cdots C_{k-1}P_{k-1}C_kQ_k$ is a directed Eulerian walk.

CASE 3: $\boldsymbol{u} \in V(Q_k)$.

Let i_{k+1} be the index of $\boldsymbol{u}$ in Q_k and set $j_{k+1} := l$. Then $\boldsymbol{u} = u_{i_{k+1}} = u_{j_{k+1}}$ and $j_k < i_{k+1} < l$. Define the following cycle and open path

$$C_{k+1} = u_{i_{k+1}}x_{i_{k+1}+1}u_{i_{k+1}+1}x_{i_{k+1}+2}\ldots u_{j_{k+1}-1}x_{j_{k+1}}u_{j_{k+1}}, \tag{2.4}$$

$$P_k = x_{j_k+1}u_{j_k+1}x_{j_k+2}u_{j_k+2}\ldots x_{i_{k+1}-1}u_{i_{k+1}-1}x_{i_{k+1}}. \tag{2.5}$$

Suppose that Q_k meets some of $C_1,\ldots,C_k$. Let α be the least index of vertices of $V(Q_k)\cap V(C_1\cup\cdots\cup C_k)$ in Q_k. Then we must have passed the step where $\ell(\boldsymbol{x}) = \alpha$. According to CASE 2, we had found a directed Eulerian sub-walk of W at $\ell(\boldsymbol{x}) = \alpha$, and could not have run into the current situation. So Q_k is vertex-disjoint from $C_1,\ldots,C_k$.

If $\varepsilon(\boldsymbol{u},\boldsymbol{x}) \neq \varepsilon(u_{i_{k+1}}, x_{i_{k+1}+1})$, then $W_0 := C_{k+1}$ is a directed circuit of Type I and is a sub-walk of W. STOP; see Figure 12.



Figure 12. Left: Q_k is disjoint from $C_1P_1\cdots C_{k-1}P_{k-1}C_k$. Right: Q_k intersects some of $P_1,\ldots,P_{k-1}$.

If $\varepsilon(\boldsymbol{u},\boldsymbol{x}) = \varepsilon(u_{i_{k+1}}, x_{i_{k+1}+1})$, then C_{k+1} is an unbalanced cycle. Since $\boldsymbol{u} \neq u_{i_1} = u_0$, the function f_W is unbalanced at $\boldsymbol{u}$. We first consider the situation that Q_k is disjoint from $P_1,\ldots,P_{k-1}$, that is,

$$V(Q_k) \cap V(P_1 \cup \cdots \cup P_{k-1}) = \emptyset.$$

There exists an edge $x' = \boldsymbol{u}u' \in \operatorname{supp} f'_W$ such that $\varepsilon(\boldsymbol{u},\boldsymbol{x}) \neq \varepsilon(\boldsymbol{u},x')$; see the left of Figure 13. Define $\ell(x') = l+1$, write $\boldsymbol{u}x'$ as $u_l x_{l+1}$, and set $W := Wu_l x'$. Assign (x',u') to $(\boldsymbol{x},\boldsymbol{u})$ and return to STEP 4 with k replaced by $k+1$.

Next we consider the situation that Q_k meets some of $P_1,\ldots,P_{k-1}$, that is,

$$V(Q_k) \cap V(P_1 \cup \cdots \cup P_{k-1}) \neq \emptyset.$$

Let γ be the greatest index of vertices of $V(Q_k) \cap V(P_1 \cup \cdots \cup P_{k-1})$ in Q_k. Let α be the index of u_γ in $P_1 \cup \cdots \cup P_{k-1}$, say, in P_r with $1 \leq r < k$. Then $j_k < \gamma < l$, $j_r < \alpha < i_{r+1}$, and $u_\alpha = u_\gamma$. Recall $\boldsymbol{u} = u_{i_{k+1}} = u_{j_{k+1}}$ and $j_{k+1} = l$, $j_k < i_{k+1} < l$. We continue with the following three subcases.



Figure 13. Left: $\varepsilon(\boldsymbol{u},\boldsymbol{x}) = \varepsilon(u_{i_{k+1}}, x_{i_{k+1}+1})$. Right: $\varepsilon(\boldsymbol{u},\boldsymbol{x}) \neq \varepsilon(u_{i_{k+1}}, x_{i_{k+1}+1})$.

Subcase 3.1: $\gamma < i_{k+1}$.

Since f_W is not conservative at $\boldsymbol{u}$, there exists an edge $x' = \boldsymbol{u}u' \in \operatorname{supp} f'_W$ such that $\varepsilon(\boldsymbol{u},\boldsymbol{x}) \neq \varepsilon(\boldsymbol{u},x')$. Define $\ell(x') = l+1$, write $\boldsymbol{u}x'$ as $u_l x_{l+1}$, and set $W := W\boldsymbol{u}x'$. Assign (x',u') to $(\boldsymbol{x},\boldsymbol{u})$ and return to STEP 4 with k replaced by $k+1$. See the right of Figure 13.

Subcase 3.2: $\gamma = i_{k+1}$.

Note that $\boldsymbol{u} = u_l = u_{i_{k+1}} = u_\gamma = u_\alpha$. Then $W_0 := C_{r+1}P_{r+1}\cdots C_kP_kC_{k+1}P'_r$ is a directed Eulerian sub-walk of W, where P'_r is a path defined by (2.3). STOP; see Figure 14.

Subcase 3.3: $\gamma > i_{k+1}$. There are two situations.



FIGURE 14. An Eulerian walk $C_{r+1}P_{r+1}\cdots C_kP_kC_{k+1}P_r'$.



FIGURE 15. $C_{r+1}P_{r+1}\cdots C_kP_kQ_{k+1}''P_r'$ is a directed Eulerian walk.

Situation 3.3.1: $\boldsymbol{u} \notin V(P_1 \cup \cdots \cup P_{k-1})$.

Notice that we must have passed CASE 1 when the variable $\boldsymbol{u} = u_\gamma$. Since we have run into the current step, we must have $\varepsilon(u_\gamma, x_\gamma) = \varepsilon(u_\alpha, x_{\alpha+1})$. Then $W_0 := C_{r+1}P_{r+1}\cdots C_kP_kQ_{k+1}''P_r'$ is a directed Eulerian sub-walk of W, where P_k is a path defined by (2.5), Q_{k+1}'' is defined by

$$Q_{k+1}'' = u_lx_lu_{l-1}x_{l-1}u_{l-2}\ldots x_{\gamma+2}u_{\gamma+1}x_{\gamma+1}(u_\gamma), \tag{2.6}$$

and P_r' is a path defined by (2.3). STOP; see Figure 15.

Situation 3.3.2: $\boldsymbol{u} \in V(P_1 \cup \cdots \cup P_{k-1})$.

It is similar to **Situation 3.3.1** that $W_0 := C_{r+1}P_{r+1}\cdots C_kP_kQ_{k+1}''P_r'$ is a directed Eulerian sub-walk of W, where P_r' is defined by (2.3) and Q_{k+1}'' is defined by (2.6). STOP; see Figure 16.



FIGURE 16. $C_{r+1}P_{r+1}\cdots C_kP_kQ_{k+1}''P_r'$ is a directed Eulerian walk.

3. Proof of Main Results

We first prove that the supports of directed Eulerian walks obtained by the *Flow Reduction Algorithm (FRA)* are Eulerian cycle-trees. Then we prove Theorem 1.2, Theorem 1.1 and Theorem 1.3. Corollaries 1.4 and 1.5 are direct consequences of Theorems 1.1–1.3.

Lemma 3.1. *The support of each directed Eulerian walk* (W, ε_W) *found by FRA is an Eulerian cycle-tree.*

Proof. Check the FRA; we see that the walk W has the following four types:

(a) $W = C_1$ is a balanced cycle, a circuit of Type I.
(b) $W = C_1C_2$, where C_1, C_2 are two unbalanced cycles having exactly one intersection vertex, a circuit of Type II.
(c) $W = C_1PC_2P^{-1}$, where C_1, C_2 are vertex-disjoint unbalanced cycles and P is a simple path intersecting C_1 exactly at the initial vertex and C_2 exactly at the terminal vertex. The support of W is a circuit of Type III.
(d) $W = C_1P_1C_2P_2 \cdots C_kP_kC_{k+1}$ with $k \geq 2$ and $C_{k+1} = C_1$, where C_i are vertex-disjoint unbalanced cycles, each P_i is a simple open path such that $C_iP_iC_{i+1}$ is a circuit of Type III and vertex-disjoint from all cycles other than C_i, C_{i+1}, the vertices and edges of P_i may appear twice in W.

We call a vertex (an edge) of $\operatorname{supp} W$ a *double vertex* (*double edge*) if it appears twice in W. A *double-edge path* is a path of $\operatorname{supp} W$ whose edges are double edges of W. A double-edge path is said to be *maximal* if it is not contained strictly in a longer double-edge path as edge subsets.

For the circuit case of (a), (b) and (c), it is clear that $\operatorname{supp} W$ is a cycle-tree. For the case (d), to see that $\operatorname{supp} W$ is a cycle-tree, we claim that every double vertex of $\operatorname{supp} W$ is a separable vertex. If so, let $\{P'_s\}$ denote the set of all maximal double-edge paths of $\operatorname{supp} W$. Then P'_s are vertex-disjoint. Remove all P'_s from $\operatorname{supp} W$ but keep their initial and terminal vertices, the leftover is a collection of blocks which are all cycles, denoted C'_r. Then $T_W = \{C'_r, P'_s\}$ is a cycle-tree.

Let u be a double vertex of $\operatorname{supp} W$. If u is the initial vertex or terminal vertex of a path P_i in W, it is clear that u is a separable vertex because the cycles C_i and C_{i+1} are separated by the initial and terminal vertices of P_i, according to the construction of W.

Let u be the internal intersection vertex of two paths P_a, P_c with $a < c$. We write $W = u_0x_1u_1x_2 \ldots u_{l-1}x_lu_l$ with $u_l = u_0$. Let u be indexed as u_α in P_a and as u_γ in P_c. Then by the intersection pattern of FRA, we have

$$\varepsilon_W(u_\gamma, x_\gamma) = \varepsilon_W(u_\alpha, x_{\alpha+1}) = -\varepsilon_W(u_\alpha, x_\alpha) = -\varepsilon_W(u_\gamma, x_{\gamma+1}).$$

It is possible that $x_\alpha u_\alpha x_{\alpha+1} = x_{\gamma+1}u_\gamma x_\gamma$. Suppose u is not a separable vertex of $\operatorname{supp} W$. This means that the closed walks

$$u_0x_1u_1 \ldots x_\alpha u_\alpha x_{\gamma+1}u_{\gamma+1} \ldots u_{l-1}x_lu_l, \quad u_\alpha x_{\alpha+1}u_{\alpha+1} \ldots u_{\gamma-1}x_\gamma u_\gamma$$

intersects at a vertex v other than u. Let v be indexed as $u_\beta \in P_b$ and $u_\delta \in P_d$, where $a \leq b \leq c$, $d \leq a$ or $d \geq c$. If $d \leq a$, then $\delta < \alpha < \beta < \gamma$. If $d \geq c$, then $\alpha < \beta < \gamma < \delta$. In both cases the intersection pattern at double vertices implies that

$$\varepsilon_W(u_\beta, x_\beta) = \varepsilon_W(u_\delta, x_{\delta+1}) = -\varepsilon_W(u_\delta, x_\delta) = -\varepsilon_W(u_\beta, x_{\beta+1}).$$

In the case of $\delta < \alpha < \beta < \gamma$, the FRA finds a directed Eulerian walk

$$W_1 = u_\delta x_{\delta+1} u_{\delta+1} \ldots u_{\alpha-1} x_\alpha u_\alpha (u_\gamma) x_\gamma u_{\gamma-1} x_{\gamma-1} \ldots u_{\beta+1} x_{\beta+1} u_\beta$$

at the step where $\ell(\boldsymbol{x}) = \gamma$. Then we shall not run RFA further at the step of $\ell(\mathbf{x}) = \gamma$ to obtain W. This is a contradiction; see FIGURE 17. In the case of $\alpha < \beta < \gamma < \delta$, the FRA finds a directed Eulerian walk



FIGURE 17. Non-separability implies decomposability.

$$W_2 = u_\alpha x_{\alpha+1} u_{\alpha+1} \ldots u_{\beta-1} x_\beta u_\beta (u_\delta) x_\delta u_{\delta-1} x_{\delta-1} \ldots u_{\gamma+1} x_{\gamma+1} u_\gamma$$

at the step where $\ell(\boldsymbol{x}) = \delta$. Then we shall not run FRA further at the step of $\ell(\boldsymbol{x}) = \delta$ to obtain W. This is a contradiction; see also FIGURE 17. So u must be a separable vertex of $\operatorname{supp} W$.

The Eulerian property follows from the following Lemma 3.3. $\square$

Lemma 3.2. *Let $W = u_0 x_1 u_1 x_2 \ldots u_{l-1} x_l u_l$ be a walk with a direction ε_W. Then*

$$\varepsilon_W(u_l, x_l) = -\sigma(W)\varepsilon_W(u_0, x_1).$$

Proof. The direction ε_W must be constructed as follows:

$$\varepsilon_W(u_i, x_i) = -\sigma(x_i)\varepsilon_W(u_{i-1}, x_i), \quad \varepsilon_W(u_i, x_{i+1}) = -\varepsilon_W(u_i, x_i), \quad 1 \le i \le l.$$

Then $\varepsilon_W(u_l, x_l)$ is determined by $\varepsilon_W(u_0, x_1)$ as $\varepsilon_W(u_l, x_l) = -\sigma(W)\varepsilon_W(u_0, x_1)$. $\square$

Lemma 3.3. *Let $T = \{C_i, P_j\}$ be a cycle-tree. If T is Eulerian, then $\sigma\big(\bigcup C_i\big) = 1$. If there exists a directed Eulerian walk (W, ε_W) on T such that the edges of C_i appear once and the edges of P_j appear twice, then T is Eulerian and $\sigma(W) = 1$.*

Proof. We may think of T as a tree having C_i as vertices, and the paths P_j and the intersection vertices between cycles as edges. Then the degree of C_i is the number of intersection vertices of C_i. Thus the number of cycles that have odd number intersection vertices must be even. Since the parity of the number of negative edges of a cycle C_i is the same as the parity of the number of intersection vertices of C_i, it follows that $\sigma\big(\bigcup C_i\big) = 1$.

To show that T is Eulerian, we proceed by induction on the number of cycles of T. Take an end-cycle C_0 of T, that is, C_0 contains exactly one intersection vertex u_0. There are two cases: (i) u_0 is the intersection of C_0 and another cycle C_1; (ii) u_0 is the intersection of C_0 and a tree-path P at its initial vertex, and in this case let C_1 be another cycle intersecting P at its terminal vertex w_0. Since the edges of C_0 appear exactly once in W, it follows that $C_0 = u_0 x_1 u_1 \ldots x_l u_l$ with $u_l = u_0$ and



FIGURE 18. End-cycle of a cycle-tree T.

is a sub-walk of W; see Figure 18. Since the edges of P appear exactly twice in W, it follows that $P = v_0y_1v_1 \ldots y_mv_m$ is a sub-path of W with $v_0 = u_0$ and $v_m = w_0$; so is P^{-1}; see Figure 18.

Now remove C_0, P from T (but keep the vertex u_l in case (i) and the vertex w_0 in case (ii)) to obtain a cycle-tree T', and from W to obtain a closed walk $W' = w_0z_1w_1 \ldots z_nw_n$ on T' with $w_n = w_0$; see Figure 18. Since (W, ε_W) is a closed directed walk, it is conservative at its initial (=terminal) vertex, then by Lemma 3.2 we have $\sigma(W) = 1$. Since (W, ε_W) is Eulerian, the orientations of the edge y_1 in P and P^{-1} are the same, so are the orientations of the edge y_m in P and P^{-1}. It follows that

$$\varepsilon_W(u_0, x_1) = \varepsilon_W(u_l, x_l) = -\varepsilon_W(v_0, y_1),$$

$$\varepsilon_W(w_0, z_1) = \varepsilon_W(w_n, z_n) = -\varepsilon_W(v_m, y_m).$$

Thus the cycle C_0 must be unbalanced by Lemma 3.2 and has odd number of intersection vertices (in fact exactly one intersection vertex); and subsequently, $\sigma(W') = -1$.

Let T'' denote the cycle-tree T' having the sign of the edge z_1 switched. Let $\varepsilon_{W'} = \varepsilon_W$ except $\varepsilon_{W'}(w_0, z_1) = -\varepsilon_W(w_0, z_1)$. Then $(W', \varepsilon_{W'})$ is a directed Eulerian walk on T''. By induction, the cycle-tree T'' is Eulerian. Note that C_1 has different parity in T and T'', and that the number of intersection vertices of C_1 are differ by one in T and T''. We see that the number of intersection vertices of C_1 is even if C_1 has positive sign, and is odd if C_1 has negative sign. So T satisfies the Parity Condition. We have proved that T is Eulerian. □

Proof OF THEOREM 1.2.

Let $T = \{C_i, P_j\}$ be a cycle-tree. We first prove the existence of the required closed walk W by induction on the number of cycles of T.

Take an end-cycle C_0 whose unique intersection vertex is u_0. Then C_0 intersects either (i) a cycle C_1 at u_0 or (ii) a tree-path P at its initial vertex u_0. In case (ii) let P intersect another cycle C_1 at its terminal vertex w_0. Write $C_0 = u_0x_1u_1 \ldots x_lu_l$ with $u_l = u_0$, and $P = v_0y_1v_1 \ldots y_mv_m$ with $v_0 = u_0$ and $v_m = w_0$. For the case (i), we remove the cycle C_0 from T but keep the vertex u_0 to obtain a cycle-tree T'. For the case (ii), we remove the cycle C_0 and the path P but keep the terminal vertex w_0 to obtain a cycle-tree T'. In both cases, T' has one fewer number of cycles and the cycle C_1 has one fewer number of intersection vertices. By induction, the vertices and edges of T' can be ordered as a required closed walk $W' = w_0z_1w_1 \ldots z_nw_n$ with $w_n = w_0$ (in the first case $w_0 = v_0 = u_0$), that is, edges on cycles of T' appear exactly once in W' and edges on tree-paths of T' appear exactly twice in W'. Then $W := C_0W'$ for the case (i) and $W := C_0PW'P^{-1}$ for the case (ii) are required closed walks constructed on T; see Figure 18.

Next let T satisfy further the Parity Condition. However, the cycle-tree T' does not satisfy the Parity Condition because of C_1. Let us switch the sign of z_1 in T' and denote by T'' the same cycle-tree after switching. Then T'' is an Eulerian cycle-tree. Let W'' denote the same closed walk W' having the sign of z_1 switched. Then by induction, there exist two opposite directions $\pm\varepsilon_{W''}$ of W'' such that $\varepsilon_{W''}$ has the same orientations on the double edges of W'', that is, $(W'', \varepsilon_{W''})$ is a directed Eulerian walk. Since z_1 has opposite signs in T' and T'', the two directions $\pm\varepsilon_{W''}$ induce two directions $\pm\varepsilon_{W'}$ on W', where $\varepsilon_{W'} = \varepsilon_{W''}$ except $\varepsilon_{W'}(w_0, z_1) = -\varepsilon_{W''}(w_0, z_1)$. Thus $(W', \varepsilon_{W'})$ is a directed walk having either a sink or source at w_0, that is, $\varepsilon_{W'}(w_0, z_1) = \varepsilon_{W'}(w_n, z_n)$. Clearly, $\varepsilon_{W'}$ has the same orientations on the double edges of W'. So $\varepsilon_{W'}$ induces an orientation $\varepsilon_{T'}$ on T' having a unique sink or source at w_0.

Let ε_P be a direction of P such that $\varepsilon_P(v_m, y_m) = -\varepsilon_{W'}(w_0, z_1) = -\varepsilon_{W'}(w_n, z_n)$, and ε_{C_0} a direction of C_0 such that $\varepsilon_{C_0}(u_0, x_1) = \varepsilon_{C_0}(u_l, x_l) = -\varepsilon_P(v_0, y_1)$. Let the direction of P^{-1} be selected to be the same as ε_P. Then the directions $\varepsilon_{C_0}, \varepsilon_P, \varepsilon_{W'}$ give rise to a direction ε_W of W so that (W, ε_W) is a directed Eulerian walk. It is clear that ε_W has the same orientations on double edges of W. So ε_W induces a direction ε_T on T.

Finally we show that the directed Eulerian walk (W, ε_W) is minimal. Let (W_1, ε_{W_1}) be a directed Eulerian walk such that the multiset $M(W_1, \varepsilon_{W_1})$ is contained in $M(W, \varepsilon_W)$. It is clear that T_1 $(= \operatorname{supp} W_1)$ is a cycle-tree and is contained in $T = \operatorname{supp} W$. Remove all maximal double-edge paths of T_1; the leftover H in T_1 after removal is a signed subgraph of T, having degree 2 or 4 (even degree) at every vertex. Then H can be decomposed into edge-disjoint cycles, which of course are some cycles of the cycle-tree T.

If H contains all cycles of T, we claim that T_1 contains all double edges of T. Suppose T has a double edge e that is not a double edge of T_1. Let P_1 be the maximal double-edge path containing e in T. Joining P_1 to T_1, there exists a unique cycle C in $T_1 \cup P_1$ such that $e \in C$. The cycle C is certainly different from all cycles of the cycle-tree T. This is a contradiction. Hence $T_1 = T$, and subsequently, $W_1 = W$.

If H does not contain all cycles of T, then the cycles of T that are not contained in H together with all maximal double-edge paths between them form some cycle-trees $T_2, \ldots, T_k$ with $k \geq 2$. Thus $T_1, T_2, \ldots, T_k$ are disjoint cycle-trees contained in T. Then there exists a double edge path P_2 in T from a cycle C' of T_1 to a cycle of T_2. Let x, y be edges of C' incident with the initial vertex u of P_2 (possibly x is a loop and $x = y$). On the one hand, since C' is a cycle of T_1 and whose edges appear exactly once in W_1, we see that $\varepsilon_{W_1}(u, x) = -\varepsilon_{W_1}(u, y)$. On the other hand, since $T = \operatorname{supp} W$ and the double-edge path P_2 is contained in T, we must have $\varepsilon_W(u, x) = \varepsilon_W(u, y)$. However, $\varepsilon_W = \varepsilon_{W_1}$ on T_1. This is a contradiction. □

Proof OF THEOREM 1.1.

Let f be a nontrivial integral flow of (Σ, ε). Consider the nonnegative integral flow $|f|$ of (Σ, ε_f), where ε_f is defined by (2.1). Applying FRA, there exists a directed Eulerian walk (W_1, ε_f) on $(\operatorname{supp}|f|, \varepsilon_f)$ such that $I_{W_1} \leq |f|$, where I_{W_1} is the indicator function of $\operatorname{supp} W_1$ defined by (1.7). According to Lemma 3.1, Lemma 3.3 and Theorem 1.2, we see that W_1 is a minimal directed Eulerian walk. If $|f| - I_{W_1} \neq 0$, then by FRA there exists a minimal Eulerian W_2 on

$(\operatorname{supp} f_{W_1}, \varepsilon_f)$ such that $I_{W_2} \le |f| - I_{W_1}$. Likewise, if $|f| - I_{W_1} - I_{W_2} \ne 0$, then by FRA there exists a minimal Eulerian walk W_3 on $(\operatorname{supp} f_{W_2}, \varepsilon_f)$ such that $I_{W_3} \le |f| - I_{W_1} - I_{W_2}$. Continue this procedure, we obtain minimal Eulerian walks $W_1, W_2, \ldots, W_k$ on $(\operatorname{supp}|f|, \varepsilon_f)$ such that $|f| = \sum_{i=1}^{k} I_{W_i}$, where I_{W_i} are the indicator functions defined by (1.7). Therefore, $[\varepsilon, \varepsilon_f] I_{W_i}$ are integral flows of (Σ, ε) and $f = \sum_{i=1}^{k} [\varepsilon, \varepsilon_f] I_{W_i}$. Note that $f_{W_i} = [\varepsilon, \varepsilon_f] I_{W_i}$ by definition (1.6). So $f = \sum_{i=1}^{k} f_{W_i}$. □

Proof OF THEOREM 1.3.

Let (W, ε_W) be a minimal directed Eulerian walk. Theorem 1.1 implies that there exist minimal directed Eulerian walks (W_i, ε_i) such that $f_W = \sum_{i=1}^{k} f_{W_i}$. Then $M(W_i, \varepsilon_i)$ are sub-multisets of $M(W, \varepsilon_W)$. Since (W, ε_W) is minimal, it follows that $k = 1$ and $M(W_1, \varepsilon_1) = M(W, \varepsilon)$, $f_W = f_{W_1}$, and W_1 is obtained from FRA. Thus $\operatorname{supp} W = \operatorname{supp} W_1$. Lemma 3.1 implies that $\operatorname{supp} W_1$ is an Eulerian cycle-tree. Hence W_1 has the following forms:

(a) $W_1 = C_1$ is a balanced cycle, a circuit of Type I.
(b) $W_1 = C_1 C_2$ is a circuit of Type II, where C_1, C_2 are two unbalanced cycles having exactly one intersection vertex.
(c) $W_1 = C_1 P C_2 P^{-1}$, where C_1, C_2 are vertex-disjoint unbalanced cycles and P is an edge-disjoint path intersecting C_1 at its initial vertex and C_2 at its terminal vertex, and $C_1 P C_2$ forms a circuit of Type III.
(d) $W_1 = C_1 P_1 C_2 P_2 \cdots C_k P_k C_{k+1}$ with $k \ge 2$ and $C_{k+1} = C_1$, where C_i are vertex-disjoint unbalanced cycles, each P_i is a simple path such that each $C_i P_i C_{i+1}$ is a circuit of Type III and vertex-disjoint from all cycles other than C_i, C_{i+1}, and P_i may intersect with patterns in Figure 1.

For the cases (a)–(c), we have $W = W_1$ up to reversing the orders of the vertex-edge sequences of C_1, C_2 and P. Thus (W, ε_W) is a directed circuit and f_W is the characteristic vector of the directed circuit. For the case (d), $W = W_1$ up to reversing the orders of the vertex-edge sequences of all P_i and the order $C_1, C_2, \ldots, C_k$, and up to reversing the orders of the vertex-edge sequences of C_i arbitrarily. Then $\operatorname{supp} W$ is an Eulerian cycle-tree. Note that each $C_i P_i C_{i+1}$ is a circuit of Type III. The function $\frac{1}{2} I_{C_i P_i C_{i+1}}$ has constant value $\frac{1}{2}$ on the cycles C_i, C_{i+1} and constant value 1 on the paths P_i. Since each C_i appears twice in two such circuits of Type III, we thus have

$$f_W = \frac{1}{2} \sum_{i=1}^{k} [\varepsilon, \varepsilon_W] I_{C_i P_i C_{i+1}}.$$

We have finished the proof for all parts of Theorem 1.3. □

REFERENCES

[1] M. Beck, T. Zaslavsky, The number of nowhere-zeo flows on graphs and signed graphs, *J. Combin. Theory Ser. B* **96** (2006), 901–918.
[2] B. Bollobás, *Modern Graph Theory*, Springer, 1998.
[3] J.A. Bondy, U.S.R. Murty, *Graph Theory*, Springer, 2008.
[4] A. Bouchet, Nowhere-zero integral flows on a bidirected graph, *J. Combin. Theory Ser. B* **34** (1983), 279–292.
[5] B. Chen, J. Wang, The flow and tension spaces and lattices of signed graphs, *European J. Combin.* **30** (2009), 263–279.

[6] B. Chen, J. Wang, Torsions formulas for signed graphs, *Discrete Appl. Math.* **158** (2010), 1148–1157.
[7] C. Godsil, G. Royle, *Algebraic Graph Theory*, Springer, 2004.
[8] F. Harary, On the notion of balance of a signed graph, *Michigan Math. J.* **2** (1953), 43–146.
[9] F. Herder, An attitudes and cognitive organization, *J. Psychology* **21** (1946),107–112.
[10] A. Khelladi, Nowhere-zero integral chains and flows in bidirected graphs, *J. Combin. Theory Ser. B* **43** (1987), 95–115.
[11] W.T. Tutte, *Graph Theory*, Encyclopedia of Math. and Its Appl., Vol **21**, Addison-Wesley, 1984.
[12] W.T. Tutte, *Introduction to the Theory of Matroids*, Elsevier, New York, 1971.
[13] W.T. Tutte, A class of abelian groups, *Candian J. Math.* **8** (1956), 13–28.
[14] T. Zaslavsky, Signed graphs, *Discrete Appl. Math.* **4** (1982), 47–74.
[15] T. Zaslavsky, Orientation of signed graphs, *European J. Combin.* **12** (1991), 361–375.

Department of Mathematics, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong
E-mail address: `mabfchen@ust.hk`

Department of Mathematics and Physics, Shenzhen Polytechnic, Shenzhen, Guangdong Province, 518088, P.R. China
E-mail address: `twojade@alumni.ust.hk`